

Dentistry Compounds

**** CALL PHARMACY FOR PRICING DETAILS ****

1-800-523-1486

1. Carsol

- Liquid
- Ingredients: Benzocaine, Dexamethasone, Polymyxin B, Bacitracin, Alcohol (pga), propylene glycol, cinnamon
- Uses: aphthous ulcer (canker sore)
- Comments: May apply 4 to 6 times daily to sore. Do not eat or drink for 30 minutes after dose.

2. Negatan

- Ingredients: Polyphenolsulfonic acid and formaldehyde solution
- Uses: aphthous ulcer (canker sore). Skin antiseptic and disinfectant.
- Comments: The prescription has to sit for 48 hours after being made prior to dispensing. May apply 4 to 6 times daily to sore. Do not eat or drink for 30 minutes after dose.

3. Viruscide Cream

- Ingredients: Alaskan Marine Red Algae – antiviral, antibacterial, and immune boosting properties
- Uses: Fever blister (cold sore) on outside of mouth
- Comments: Apply to the affected area every 30 minutes while awake for the first 24 hours, then apply every 1 to 2 hours. Best results obtained if applied at first symptom.

4. MD Anderson Mouth Wash or Magic Mouth Wash

- Ingredients: Tetracycline 500mg, Nystatin 15ml, Hydrocortisone 100mg, Benadryl qs to 240ml.
- Uses: mouth pain, canker sores, chemotherapy/radiation-induced mucositis.
- Comments: Use every 4 to 6 hours with instructions to hold in the mouth for one to two minutes then spit out or swallow. Patients should be instructed not to eat or drink for 30 minutes after use. We can provide

any combination of ingredients which includes: Tetracycline, Nystatin, Hydrocortisone, Benadryl, Viscous Lidocaine, Maalox.

5. Lidocaine 10%/Prilocaine 10%/Tetracaine 4% Gel

- Uses: local anesthetic gel. This gel is used locally prior to injections so patients do not hardly feel these painful injections.
- Comment: Apply a small amount locally prior to injection. Numbing effect enhances danger of aspiration. Don't eat, drink, or chew gum while anesthetized.

6. Dyclonine HCL 1% Oral Dental Solution

- Uses: local anesthetic with bacteriocidal and fungicidal properties. It is used for oral pain and also used as a mouthwash or gargle in suppressing the gag reflex.
- Comment: Rinse 30ml for no more than 1 minute and expectorate. Numbing effect enhances danger of aspiration. Don't eat, drink, or chew gum while anesthetized. Use in patients with "caine" sensitivity.

7. Pilocarpine

- Uses: dry mouth, Sjogrens syndrome. Cholinergic agonists work by increasing the amount of saliva in the mouth.
- Comments: side effect of excessive sweating. Do not use anticholinergic medications with pilocarpine.

8. Benzocaine 16.7% Dry Socket Paste

- Uses: relief of pain of mouth and gum sores, mouth pain, canker sores, irritation of the mouth.
- Comments: Apply to the socket with gauze up to four times daily.

9. Tretinoin 0.1%/Clobetasol Propionate 0.05% Oral Adhesive Paste

- Uses: inflammation and irritation of oral dermatoses
- Comments: Apply topically three to four times daily and frequency should be tapered as symptoms improve.