

Pediatric Compounds

**** CALL PHARMACY FOR PRICING DETAILS ****

1-800-523-1486

Diaper Rash Compounds:

- 1. Buttocks Ointment:** Contains (Peruvian Balsam, Castor Oil, Boric Acid, Zinc Oxide, Mineral Oil, Petrolatum)
 - For diaper rash with yeast add: Nystatin 100,000 Units/gm
 - For really irritated/red/itchy add: Hydrocortisone 1% or Triamcinolone 0.025%
 - 2. Cholestyramine 5% Diaper Rash Ointment:** mix with Aquaphor.
 - This ointment is great for diarrhea as the cholestyramine binds to the bile acids and prevents it from burning the baby's bottom.
 -
 - 3. Happy Hiney:** Equal parts of (Aquaphor/Monistat(miconazole)/1% hydrocortisone/Maalox)
 - This ointment is great and is a popular diaper rash ointment in TN.
- ❖ **Uses:** Diaper Rash
 - ❖ **Precautions/warnings:** Hypersensitivity to any ingredients
 - ❖ **Comments:** These are great **"GO TO"** diaper rash treatments that are both effective and affordable for the parents, and makes their little baby's booty healthy & happy!

Nausea/Vomiting:

- 1. Ondansetron 2mg/0.1ml, 4mg/0.1ml or 8mg/0.1ml in Lipoderm**
 - 2. Promethazine 25mg/0.1ml or 12.5mg/0.1ml in Lipoderm**
- ❖ **Uses:** Nausea/vomiting
 - ❖ **Precautions/warnings:** Do not administer with apomorphine EVER! (apomorphine not approved for children)
 - ❖ **Comments:** Use acutely to stop n/v in patients where dehydration is a concern. Can also be used if giving another medication that may cause n/v. This is an excellent dosage form for n/v since it does not have to be given orally to children who are vomiting already.

Acid Reducers:

- 1. Ranitidine 15mg/ml oral suspension**
 - 2. Omeprazole 2mg/ml oral suspension**
 - 3. Lansoprazole 3mg/ml oral suspension**
- ❖ **Uses:** GERD
 - ❖ **Precautions/warnings:** Hypersensitivity to active ingredients.
 - ❖ **Comments:** OUR STUFF TASTES BETTER! The manufactured drugs are either unavailable or taste awful. Our suspensions are dye-free. We can compound suspensions for 14 days supply at a time and all must be kept in the refrigerator.

Acne and Skin condition Solutions:

1. Benzoyl Peroxide 15% or 20%/ Erythromycin 4% (mix in versa base)

- ❖ **Uses:** Acne (benzoyl peroxide is an oxidizing agent with antibacterial properties, erythromycin, antibacterial)
- ❖ **Precautions/warnings:** Hypersensitivity to either ingredient.
- ❖ **Comments:** Patients should use a sunscreen while using this medication. May be used twice daily or at bedtime. If excessive dryness occurs, decrease amount of product used and frequency. Use a moisturizer.

2. Tretinoin 0.025% or 0.05% can be with or without Hydroquinone 4% cream

- ❖ **Uses:** For improving or removing dark spots, blemishes & dead skin cells. It may also be used for other conditions as determined by your doctor.
- ❖ **Precautions/warnings:** hypersensitivity to any ingredient
- ❖ **Comments:** Patients should not be in the sun without the use of sunscreen while using this medication. May be used twice daily or at bedtime. If excessive dryness occurs, decrease amount of product used and frequency. Use a moisturizer.

3. Newman's Nipple Ointment (Mupirocin Ointment 2%/Betamethasone Dipropionate 7.5mg/Miconazole 300mg in a Vaseline base)

- ❖ **Uses:** Soothe nipples during breastfeeding/prevention of thrush .
- ❖ **Precautions/warnings:** hypersensitivity to any ingredient by mother or baby
- ❖ **Comments:** apply after every feeding, and there is no need to wash off before feeding.

4. Hydrocortisone 2.5%/Aquaphor

- ❖ **Uses:** Eczema/dermatitis
- ❖ **Precautions/warnings:** Hypersensitivity to ingredients
- ❖ **Comments:** Apply once daily after bathing to affected area.